



SURF1 gene

SURF1, cytochrome c oxidase assembly factor

Normal Function

The *SURF1* gene provides instructions for making a protein that is important in oxidative phosphorylation, the process by which the energy from food is converted into a form cells can use. Oxidative phosphorylation involves a series of reactions that take place through several different protein complexes. The SURF1 protein aids in the correct assembly of one of the protein complexes, or enzymes, involved in oxidative phosphorylation called complex IV.

Complex IV, also known as cytochrome c oxidase or COX, accepts negatively charged particles (electrons) from earlier steps in oxidative phosphorylation. In addition, the enzyme accepts positively charged particles (protons) from inside the mitochondrion. Using the electrons and protons, the COX enzyme performs a chemical reaction that converts oxygen to water. The enzyme also transfers additional protons across the specialized membrane inside the mitochondrion. These processes create energy that is used to generate adenosine triphosphate (ATP), the cell's main energy source.

Health Conditions Related to Genetic Changes

cytochrome c oxidase deficiency

Leigh syndrome

More than 80 different *SURF1* gene mutations have been identified in people with Leigh syndrome, a progressive brain disorder that usually appears in infancy or early childhood. Affected children may experience delayed development, muscle weakness, problems with movement, or difficulty breathing.

Approximately 10 to 15 percent of people with Leigh syndrome have a mutation in the *SURF1* gene. Most *SURF1* gene mutations result in an abnormally short protein. Other mutations replace a single protein building block in the SURF1 protein. The mutated proteins are broken down in the cell, which results in the absence of SURF1 protein. Lack of SURF1 protein hinders the proper formation of the COX complex. As a result, COX enzyme activity is severely reduced, which leads to impaired oxidative phosphorylation.


Although the exact mechanism is unclear, researchers believe that impaired oxidative phosphorylation can lead to cell death because of decreased energy available in the cell. Certain tissues that require large amounts of energy, such as the brain, muscles,

and heart, seem especially sensitive to decreases in cellular energy. Cell death in the brain likely causes the characteristic changes in the brain seen in Leigh syndrome, which contribute to the signs and symptoms of the condition. Cell death in other sensitive tissues may also contribute to the features of Leigh syndrome.

Chromosomal Location

Cytogenetic Location: 9q34.2, which is the long (q) arm of chromosome 9 at position 34.2

Molecular Location: base pairs 133,351,805 to 133,356,485 on chromosome 9 (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

Other Names for This Gene

- SURF-1
- SURF1_HUMAN
- surfeit 1
- surfeit locus protein 1

Additional Information & Resources

Educational Resources

- Biochemistry (5th Edition, 2002): Cytochrome c Oxidase Catalyzes the Reduction of Molecular Oxygen to Water
<https://www.ncbi.nlm.nih.gov/books/NBK22505/#A2517>
- Biochemistry (5th Edition, 2002): The Respiratory Chain Consists of Four Complexes: Three Proton Pumps and a Physical Link to the Citric Acid Cycle
<https://www.ncbi.nlm.nih.gov/books/NBK22505/>

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28%28SURF1%5BTIAB%5D%29+OR+%28surfeit+1%5BTIAB%5D%29%29+OR+%28surfeit+locus+protein+1%5BTIAB%5D%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+2160+days%22%5Bdp%5D>

OMIM

- SURFEIT 1
<http://omim.org/entry/185620>

Research Resources

- Atlas of Genetics and Cytogenetics in Oncology and Haematology
http://atlasgeneticsoncology.org/Genes/GC_SURF1.html
- ClinVar
<https://www.ncbi.nlm.nih.gov/clinvar?term=SURF1%5Bgene%5D>
- HGNC Gene Family: Mitochondrial respiratory chain complex assembly factors
<http://www.genenames.org/cgi-bin/genefamilies/set/645>
- HGNC Gene Symbol Report
http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/hgnc_data.php&hgnc_id=11474
- NCBI Gene
<https://www.ncbi.nlm.nih.gov/gene/6834>
- UniProt
<http://www.uniprot.org/uniprot/Q15526>

Sources for This Summary

- Finsterer J. Leigh and Leigh-like syndrome in children and adults. *Pediatr Neurol.* 2008 Oct;39(4): 223-35. doi: 10.1016/j.pediatrneurol.2008.07.013. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/18805359>
- Lee IC, El-Hattab AW, Wang J, Li FY, Weng SW, Craigen WJ, Wong LJ. SURF1-associated Leigh syndrome: a case series and novel mutations. *Hum Mutat.* 2012 Aug;33(8):1192-200. doi: 10.1002/humu.22095. Epub 2012 Apr 30.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/22488715>
- Pecina P, Capková M, Chowdhury SK, Drahota Z, Dubot A, Vojtísková A, Hansíková H, Houst'ková H, Zeman J, Godinot C, Houstek J. Functional alteration of cytochrome c oxidase by SURF1 mutations in Leigh syndrome. *Biochim Biophys Acta.* 2003 Sep 1;1639(1):53-63.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/12943968>

- Péquignot MO, Dey R, Zeviani M, Tiranti V, Godinot C, Poyau A, Sue C, Di Mauro S, Abitbol M, Marsac C. Mutations in the SURF1 gene associated with Leigh syndrome and cytochrome C oxidase deficiency. Hum Mutat. 2001 May;17(5):374-81.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/11317352>
 - OMIM: SURFEIT 1
<http://omim.org/entry/185620>
 - Yao J, Shoubridge EA. Expression and functional analysis of SURF1 in Leigh syndrome patients with cytochrome c oxidase deficiency. Hum Mol Genet. 1999 Dec;8(13):2541-9.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/10556303>
-

Reprinted from Genetics Home Reference:
<https://ghr.nlm.nih.gov/gene/SURF1>

Reviewed: June 2016
Published: March 21, 2017

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services